

Evidence Against the Sciama Model of Radiative Decay of Massive Neutrinos ¹

Stuart Bowyer, Eric J. Korpela, Jerry Edelstein, and Michael Lampton
Space Sciences Laboratory, University of California, Berkeley, CA 94720-7450

Carmen Morales, Juan Pérez-Mercader, José F. Gómez ², and Joaquín Traperó³
Laboratorio Astrofísica Espacial y Física Fundamental, INTA. Apdo. Correos 50727, 28080
Madrid, Spain

ABSTRACT

We report on spectral observations of the night sky in the band around 900 Å where the emission line in the Sciama model of radiatively decaying massive neutrinos would be present. The data were obtained with a high resolution, high sensitivity spectrometer flown on the Spanish MINISAT satellite. The observed emission is far less intense than that expected in the Sciama model.

Subject headings: cosmology: diffuse radiation — elementary particles — ultraviolet: general

1. Introduction

Relic neutrinos, if massive, could contribute significantly to the density of the universe, and if appropriately concentrated, could explain puzzling characteristics of luminous matter in galaxies. Melott (1984) suggested that if these particles were radiatively decaying, they could be responsible for the sharp hydrogen ionization edges seen in many galaxies and that this decay would not violate existing observational data if the decay energy was somewhat greater than 13 eV and the lifetime for decay was about 10^{24} s. In a subsequent paper,

¹Based on the development and utilization of the Espectrógrafo Ultravioleta de Radiación Difusa, a collaboration of the Spanish Instituto Nacional de Técnica Aeroespacial and the Center for EUV Astrophysics, University of California, Berkeley

²Also at Instituto de Astrofísica de Andalucía, CSIC, Apdo Correos 3004, E-18080, Granada, Spain

³Present address: Universidad SEK, Cardenal Zúñiga S/n, Segovia, Spain

Melott et al. 1988 showed this idea was consistent with observations of star formation, galaxy formation and morphology, and other phenomena. Subsequently, Sciama and collaborators in an extensive set of papers (Sciama 1990, 1993, 1995, 1997a, 1997b, 1998, Sciama et al. 1993) showed that if the decay lifetime was an order of magnitude less than that suggested by Melott his theory could explain a large number of otherwise puzzling astronomical phenomena, including the ionization state of the intergalactic medium and the anomalous ionization of the interstellar medium (ISM) in our own Milky Way Galaxy. Although massive neutrinos cannot be contemplated within the framework of the standard model of particle physics, they can be accommodated in the supersymmetric extensions of the standard model, especially if R-parity is broken (cf. Gato et al. 1985, Bowyer et al. , 1995). Recent observational and experimental results suggest they do, in fact, have mass (Fukuda et al. 1998a,b and Athanassopoulos et al. 1998a,b).

A number of searches have been made for evidence of radiatively decaying massive neutrinos in clusters of galaxies. Davidsen et al. (1991) severely constrained the parameter space available for these particles through observations of the cluster of galaxies Abell 665, and Fabian et al. 1991 obtained similar results from a study of the cluster of galaxies surrounding the quasar 3C263. However, Sciama et al. (1993) and Bowyer et al. (1995) have shown that these observations do not rule out the Sciama scenario.

An all-pervading neutrino flux in the Galaxy at a wavelength near the ionization limit of hydrogen would be difficult to observe because of absorption by the ISM. However, Bowyer et al. (1995) pointed out that this flux would be observable from Earth orbit in several well-defined directions where the density of the ISM is extremely low. An observational complexity which could complicate these measurements is emission from an upper atmosphere oxygen recombination feature at 911 Å (Chakrabarti et al. 1983).

In this paper we report results of spectral observations made in the region ≤ 912 Å where the radiation in the Sciama scenario would be present, and compare the data obtained with the flux expected.

2. Observations

The observations were made with an extreme ultraviolet spectrometer covering the band-pass from 350-1100 Å which was specifically designed for studies of diffuse emission. The instrument (the Espectrógrafo Ultravioleta extremo para la Radiación Difusa, EURD) is capable of providing measurements of the diffuse UV background which are more than 100 times more sensitive than existing measurements in this band-pass, with a spectral

resolution of about $6\text{\AA}$. The instrument is described in detail by Bowyer et al. 1997.

The instrument was flown onboard the Spanish MINISAT-01 satellite launched on April 21, 1997. The spacecraft is in a retrograde orbit with an inclination of 151° and is at an altitude of 575 km . The spectrometer continuously views the anti-Sun direction. Details of the spacecraft and the EURD observational parameters are provided in Morales et al. 1996.

We examined EURD data in the 890 to $915\text{\AA}$ bandpass in an attempt to detect the emission which would be present if the Sciama scenario was operative. Data from the spectrometer were typically collected over the entire night-time portion of the orbit. Higher count rates are always experienced at spacecraft sunrise and sunset due to geocoronal effects, but deep night intensities are typically constant and low. For the search for radiation from the Sciama scenario, we sorted the data to exclude all sunrise and sunset data and all other data associated with high backgrounds. Given the low in-flight counting rate and the absolute fixed electronics dead time of $100\text{ }\mu\text{s}$ per photon, dead time corrections were about 1% and were therefore ignored.

The EURD spectrograph employs a number of vetoes to reduce unwanted background and to permit evaluation of those background events which cannot be otherwise eliminated (Bowyer et al. 1997). The detector is surrounded by an anti-coincidence shield and all counts triggering this shield (about 20 percent) are rejected. Remaining internal background components include charged particles that are missed by the anti-coincidence system, Compton scattered γ -rays, and radioactivity within the detector and in the spacecraft. An additional background is produced by photons scattered by the grating onto the detector. This scattered emission is mostly a continuum arising from the wings of the zero and first order of the hydrogen Lyman-alpha line whose peaks were designed to fall beyond the ends of the detector.

The entrance aperture of the instrument has a filter wheel with three positions: Open, Closed, and a MgF_2 filter. The Open position provides spectral data plus backgrounds. The Closed position gives an estimate of the internal background, and the MgF_2 filter position gives an estimate of the scattered radiation. Observations were carried out sequentially with each of these apertures; the complete cycle time was 90 s.

We corrected the deep night spectral data for backgrounds using the MgF_2 and Closed apertures. We summed the background corrected data in the 890 to $915\text{\AA}$ band as a function of time. We included data to $915\text{\AA}$ to assure all counts shortward of $912\text{\AA}$ were included in the sample given the spectral resolution of the instrument. In some neutrino decay scenarios, two lines will be produced whose relative intensities are uncertain.




Fig. 1.— The observed spectrum of the region around 911 Å where r adiation in the Sciamia scenario is expected. The solid line shows a straight line interpolation of the spectrum from about 60 hours of shutter open observations during a period when emission from the oxygen recombination feature was more pronounced. The dashed line shows the expected hydrogen Lyman series lines with an electron temperature of 0.4eV folded with the instrument resolution and fit to the average intensity of the Lyman series lines during this period.

However, the sum of both of these lines is the key parameter to be measured, and in the Sciamia scenario these lines will be separated by 0.2 eV, or 13 Å at 900 Å. Hence the flux from both these lines will be included in the data reported here. For this study we utilized data obtained between 18 June 1997 and 29 June 1998. Data were regularly obtained over most of this period, with occasional gaps because of spacecraft problems or instrument shutdowns. Data were summed over 10 day intervals, providing typically about 3500 counts, to obtain good counting statistics.

Unfortunately in regards to our search for the Sciamia line, oxygen recombination radiation was substantial at the altitude of the MINISAT satellite even in the anti-Sun view direction. A spectrum of the radiation detected around 912 Å is shown in Fig. 1. This spectrum shows a profile that is consistent with the line shape obtained by Feldman et al. 1992 given the resolution of this instrument. Just longward of 912 Å the spectrum is

dominated by the Lyman series lines of geocoronal hydrogen (López-Moreno et al. 1998). Both the oxygen recombination feature and the Lyman series of hydrogen vary in time; the data shown in Fig. 1 are from a period when the oxygen recombination radiation was more pronounced.

We determined the EURD counts-to-flux conversion factor in the region around 800 Å using an in-flight calibration strategy based on simultaneous EUV observations of the Moon with EUVE and EURD (Flynn et al. 1998), and, longward of 912 Å, to fits to stellar spectra (Morales et al. , in progress). It is estimated that this calibration is good to $\pm 20\%$ in the band around 912 Å because of the quality of the fit to stellar spectra. This in-flight calibration yields a conversion of 6.5×10^4 ph cm $^{-2}$ str $^{-1}$ per count at 912 Å; this is within a factor of three of the preflight calibration (Bowyer et al. 1997). This difference is easily understood as being the result of the degradation of the detector photocathode during the almost 2 year time period between the laboratory calibration and in-orbit operation. The resulting fluxes are shown in Fig. 2. These fluxes are the total fluxes obtained in this bandpass, uncorrected for any Lyman series emission as seen in Fig. 1.

The expected emission in the Sciamia scenario can best be considered in two parts. The first is produced in the Local Interstellar Cloud (LIC) which surrounds the Sun; this emission is intermixed with absorption. The second component is emission from beyond the LIC which is absorbed by this cloud.

Formally, the emission is given by the relation:

$$I(l) = B + \frac{R_{\text{prod}}}{4\pi n_o \sigma} (1 - \exp[-n_o \sigma d_{\text{cl}}(l)]) + \frac{R_{\text{prod}} [d_e(l) - d_{\text{cl}}(l)]}{4\pi} \exp[-n_o \sigma d_{\text{cl}}(l)] \quad (1)$$

where we have included a background, B (which could be due to anything, but is mostly due to oxygen recombination radiation); R_{prod} is the photon production rate; n_o is the density of the LIC; σ is the effective ISM cross section for absorption (Rumph et al. 1994); d_{cl} is the distance to the cloud edge; and d_e is the distance to the edge of the neutral free region. The symbol l indicates variation with ecliptic longitude. The most recent (small) revision of the theory (Sciamia 1998) requires a photon production rate of $2 \pm 1 \times 10^{-16}$ s $^{-1}$ cm $^{-3}$.

We have used the model of Redfield and Linsky (1999) for data on the LIC. This is a three dimensional model which is based on ISM absorption features in the spectra of nearby stars obtained with HST, EUVE, and ground based telescopes. Minimum hydrogen columns in the plane of the ecliptic in this model are $\sim 2.5 \times 10^{16}$ cm $^{-2}$, maximum columns are $\sim 2.5 \times 10^{18}$ cm $^{-2}$.

In the region beyond the LIC, Welsh et al. 1998 used high resolution optical




Fig. 2.— The expected signature from the radiative decay of massive neutrinos as predicted by the Sciamia scenario is shown as a dashed line. This signature depends upon ecliptic longitude because of absorption by the LIC and the size of the open region beyond. The integrated intensity of the observed emission from 890 to 915 Å is also shown. Each data point is a sum of 10 days of deep night data.

spectroscopy to determine the amount of ISM sodium in the line of sight to stars within 300 pc of the Sun. They found that the ISM is essentially free of neutral gas out to more than 70 pc in most directions. Sfeir et al. 1999 have obtained an extensive set of sodium absorption data and have modeled the extent of this ionized region, or Local Bubble. We have used the $N(\text{H}) = 1 \times 10^{19} \text{ cm}^{-2}$ contour of their model, where the ionized region of the Local Bubble abruptly ends, as the limit to the region from which the Sciamia line could be detected. This contour is typically at 100 pc in the plane of the ecliptic. We have incorporated these results in Eqn. 1, and we show the expected emission in the plane of the ecliptic for the Sciamia scenario in Fig. 2.

3. Discussion and Conclusions

The geocoronal oxygen background is obvious in the data shown in Fig. 2, but in those view directions in which the absorption by the LIC is small because of the Sun’s location within the cloud, the flux from radiatively decaying neutrinos should be far more intense than the oxygen emission. It is obvious by inspection that the emission predicted by the Sciamia theory is not present.

We have fit our data shown in Fig. 2 to a model described by Eqn. 1 in which we treat the background B and the photon production rate R_{prod} as parameters. Our best fit value for B is $2200 \text{ ph s}^{-1} \text{ cm}^{-2} \text{ str}^{-1}$. Our best fit for R_{prod} is consistent with zero and has a 95% confidence upper limit of $0.6 \times 10^{-16} \text{ s}^{-1} \text{ cm}^{-3}$, which is one third of the production rate required by the theory.

The EURD data appear to be completely incompatible with the Sciamia model of radiatively decaying massive neutrinos. We believe that the only parameters in this study that could be challenged, in principle, are the conversion factor from observed EURD counts-to-flux, and the LIC model. In evaluating the calibration issue, we note that while the most accurate conversion factor can be derived from stellar spectra, this result requires substantial justification (to be discussed elsewhere) and is not necessary for this work. The EURD in-flight calibration is firmly established to within a factor of two through our in-flight observations of the intensity of the geocoronal hydrogen Lyman lines. The counts-to-flux conversion using these in-flight results would have to be incorrect by more than a factor of five to reduce the predicted emission in the Sciamia scenario to the level of the background shown in Fig. 2 if all the uncertainties are added in their worst directions. We can think of no way that this could be realized. The other factor that could be challenged, the LIC model, would have to be incorrect by a factor of twenty to reduce the observed flux to the level of the background shown in Fig. 2. This possibility is considered to be extremely unlikely (J. Linsky, private communication).

Although we believe our data rule out the Sciamia model of radiatively decaying neutrinos, we note that we cannot exclude the earlier model of Melott with its longer lifetime. In this respect, it is intriguing to note that we do observe a faint line at $\sim 710\text{\AA}$ in long integrations with the EURD instrument which we have not been able to identify as either an upper atmospheric airglow line or as emission from the interstellar medium. (Bowyer et al. , in progress)

4. Acknowledgements

We wish to acknowledge many useful discussions with Dennis Sciama. We thank Jeff Linsky and Seth Redfield for access to their model before publication, and Seth Redfield for help in utilizing this model. We than Daphne Sfeir for access to her model before publication and her help in utilizing this model. The authors wish to thank J. Cobb for devising and implementing complex data processing programs which convert the spacecraft data to forms amenable to scientific analysis. Partial support for the development of the EURD instrument was provided by NASA grant NGR 05-003-450 and INTA grant IGE 490056.

When the NASA funds were withdrawn from the instrument development at Berkeley by Ed Weiler, the instrument was completed with funds provided by S. Bowyer. The UCB analysis and interpretation is carried out through the volunteer efforts of S. Bowyer, J. Cobb, J. Edelstein, E. Korpela, and M. Lampton. The work by C. Morales and J. Trapero is supported in part by DGCYT grant PB94-0007. J.F. Gómez is supported in part by DGCYT grant PB95-0066 and Junta de Andalucia (Spain). J. Pérez-Mercader is supported by funds provided by the Spanish ministries of Education and Defense.

REFERENCES

- Athanassopoulos, C. et al. 1998a, Phys. Rev. Lett., 81, 1774
- Athanassopoulos, C., et al. 1998b, Phys. Rev. B., 58, 2489
- Bowyer, S., Lampton, M., Peltoniemi, J., & Roos, M. 1995, Phys. Rev. D, B25, 3214
- Bowyer, S., Edelstein, J., & Lampton, M. 1997, ApJ, 485, 52.
- Chakrabarti, S., Paresce, F., Bowyer, S., & Kimble, R. 1983, JGR, 88, 4898.
- Davidson, A., Kriss, G., Ferguson, H., Blair, W., Bowers, C., & Kimble, R. 1991, Nature 351, 128.
- Fabian, A., Naylor, T. & Sciama, D. 1991, MNRAS, 249, 21.
- Feldman, P., Davidson, A., Blair, W., Bowers, C., Durrance, S., Kriss, G., Ferguson, H., Kimble, R., & Long, K. 1992 Geophys. Res. Let., 19, 453.
- Flynn, B. C., Vallergera, J. V., Gladstone, G. R., & Edelstein, J. 1998, Geophys. Res. Let., 25, 3253.

- Fukuda, Y., et al. 1998a, Phys Lett B 433, 9.
- Fukuda, Y., et al. 1998b, Phys Rev Lett 81, 1562
- Gato, B., León, J., Pérez-Mercader, J., & Quirós, M. 1985, Nucl. Phys. B., 260, 203
- López-Moreno, J.J., Morales, C., Gómez, J. F., Trapero, J., Bowyer, S., Edelstein, J., Lampton, M., & Korpela, E., 1998 Geophys. Res. Let., 25, 2937.
- Melott, A. 1984 Soviet Astronomy, 28, 478.
- Melott, A., McKay, D., & Ralston, J., 1988, ApJ, 324, L43.
- Morales, C., Bowyer, S., Gómez, J. F., Trapero, J., Ulibarrena, M., Gutiérrez, F., López-Moreno, J.J., Pérez-Mercader, J., Lampton, M., & Edelstein, J. 1996, International Conference on Small Satellites: Missions and Technology (INTA: Madrid, Spain), 61.
- Redfield, S., & Linsky, J., submitted to ApJ.
- Rumph, T., Bowyer, S., & Vennes, S. 1994, AJ, 107, 2108
- Sciama, D. 1990, ApJ, 364, 549.
- Sciama, D. 1993, ApJ, 409, L25.
- Sciama, D. 1995, ApJ, 448, 667.
- Sciama, D. 1997a, ApJ, 488, 234.
- Sciama, D. 1997b, MNRAS, 289, 945.
- Sciama, D. 1998, A&A, 335, 12.
- Sciama, D., Persic, M., & Salucci, P. 1993, PASP, 105, 102.
- Sfeir, D., Lallement, R., Crifo, F., & Welsh B. 1999, A&A in press.
- Welsh, B., Crifo, F., & Lallement, R., A&A. 333, 101, 1998